

NATURE  
HAS A WAY  
OF PROTECTING  
ITS OWN.



So does  
**NITROSTAT<sup>®</sup>**  
(nitroglycerin tablets, USP)

NITROSTAT, like nature, has built-in protection—protection against degradation of nitroglycerin. In fact, NITROSTAT is more stable than conventional nitroglycerin tablets under all conditions of temperature and humidity tested.

But that's not all. Average assay still shows better than 100% of label claim for at least 60 months after manufacture of NITROSTAT. Because of this stability NITROSTAT has a five-year expiration date.\*

What's more, only NITROSTAT Tablets are available in convenient packages of 4 x 25 tablets. So emergency angina relief can be ready when needed in four different locations. "Because of its rapid action, long-established efficacy, and low cost, nitroglycerin is the most useful drug among the organic nitrates that can be given sublingually."<sup>1</sup> NITROSTAT... useful alone or as concomitant therapy with long-acting nitrates or propranolol.

\*Five-year expiration date when stored at recommended storage conditions in an unopened bottle. Data on file, Medical Affairs Dept., Parke-Davis.

REFERENCE: 1. Needleman P, Johnson EM Jr. Vasodilators and the treatment of angina, in Gilman AG, Goodman LS, Gilman A (eds): Goodman and Gilman's The Pharmacological Basis of Therapeutics, ed 6. New York, Macmillan Publishing Co, Inc, 1980, pp 819-83.

**Please see next page for brief summary of prescribing information**

**PARKE-DAVIS**

Division of Warner-Lambert Company  
Morris Plains, New Jersey 07950

## **NITROSTAT®**

(nitroglycerin tablets, USP)

**Before prescribing, please see full prescribing information.**

**A Brief Summary follows:**

### **INDICATIONS**

Nitroglycerin is indicated for the prophylaxis, treatment and management of patients with angina pectoris.

### **CONTRAINDICATIONS**

Sublingual nitroglycerin therapy is contraindicated in patients with early myocardial infarction, severe anemia, increased intracranial pressure, and those with a known hypersensitivity to nitroglycerin.

### **PRECAUTIONS**

Only the smallest dose required for effective relief of the acute anginal attack should be used. Excessive use may lead to the development of tolerance. Nitrostat tablets are intended for sublingual or buccal administration and should not be swallowed. The drug should be discontinued if blurring of vision or drying of the mouth occurs. Excessive dosage of nitroglycerin may produce severe headaches.

### **ADVERSE REACTIONS**

Transient headache may occur immediately after use. Vertigo, weakness, palpitation, and other manifestations of postural hypotension may develop occasionally, particularly in erect, immobile patients. Syncope due to nitrate vasodilation has been reported. Alcohol may accentuate the cerebral ischemia symptoms.

### **METHOD OF ADMINISTRATION**

One tablet should be dissolved under the tongue or in the buccal pouch at the first sign of an acute anginal attack. The dose may be repeated approximately every five minutes until relief is obtained. Nitrostat may be used prophylactically five to ten minutes prior to engaging in activities which might precipitate an acute attack.

### **HOW SUPPLIED**

Nitrostat is supplied in four strengths in bottles containing 100 tablets each, with color-coded labels, and in color-coded Patient Convenience Packages of four bottles of 25 tablets each.

0.15 mg (1/400 grain):

N 0071-0568-24—Bottle of 100 tablets;

N 0071-0568-13—Convenience Package.

0.3 mg (1/200 grain):

N 0071-0569-24—Bottle of 100 tablets;

N 0071-0569-13—Convenience Package.

0.4 mg (1/150 grain):

N 0071-0570-24—Bottle of 100 tablets;

N 0071-0570-13—Convenience Package.

0.6 mg (1/100 grain):

N 0071-0571-24—Bottle of 100 tablets;

N 0071-0571-13—Convenience Package.

0568G060

## **PARKE-DAVIS**

Div of Warner-Lambert Co  
Morris Plains, NJ 07950



IN CERTAIN DYSLIPIDEMIAS

*A Desirable Balance of Efficacy*

**LOPID<sup>®</sup>** 300 mg  
*(gemfibrozil capsules, USP)*

Lopid is indicated for treatment of adult patients with very high serum triglyceride levels in hyperlipoproteinemia other than type I who present a risk of pancreatitis and who do not respond adequately to diet.

© 1987 Warner-Lambert Company

**Now a  
nitroglycerin  
system that  
delivers...**



Return  
to a natural cardiac rhythm

SUSTAINED RELEASE  
**Procan<sup>®</sup> SR**  
(PROCAINAMIDE  
HYDROCHLORIDE TABLETS)

250-mg tablet/500-mg scored tablet/750-mg scored tablet  
1000-mg scored tablet



Enhances the value of oral procainamide  
for selected cardiac dysrhythmias

With modern sustained-release Procan SR, short duration of effect no longer impedes the usefulness of procainamide for maintenance therapy. A standard medication now assumes greater importance, both in terms of convenience and patient compliance.

---

Administered half as often as conventional oral procainamide

---

Wide versatility in cardiac conditions

---

Can be used concomitantly with digoxin

---

Ideal maintenance therapy

---

Procan SR...  
the original sustained-release oral procainamide